Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Modification of Heat-Related Mortality in an Elderly Urban Population by Vegetation (Urban Green) and Proximity to Water (Urban Blue): Evidence from Lisbon, Portugal

Katrin Burkart, Fred Meier, Alexandra Schneider, Susanne Breitner, Paulo Canário, Maria João Alcoforado, Dieter Scherer, and Wilfried Endlicher

Table of Contents

Table S1: Average values of (equivalent) temperature, air pollution and death counts for the elderly above the age of 65 years

Table S2: Socioeconomic and sociospatial information on parish level

Figure S1: Difference in mortality increase among those > 65 years of age with a 1°C increase in UTCI above the 95th (a) and 99th (b) percentiles lags 0-2 for nighttime LST quartiles with 95% confidence intervals. GAMs allowing for interaction between UTCI and different nighttime LST classes were adjusted for long-term and seasonal trend (6 df per year), daily averages of O₃ and PM₁₀ (lag 0-1), percent of parish population >65 years of age, building density, proportion of college graduates and percentage of population receiving social benefits. Nighttime LST quartiles were 16.6°C, 17.7°C and 18.1°C.

Figure S2: Difference in mortality increase among those > 65 years of age with a 1°C increase in UTCI above the 95th (a) and 99th (b) percentiles lags 0-2 for daytime LST quartiles with 95% confidence intervals. GAMs allowing for interaction between UTCI and different daytime LST classes were adjusted for long-term and seasonal trend (6 df per year), daily averages of O_3 and PM_{10} (lag 0-1), percent of parish population >65 years of age,

building density, proportion of college graduates and percentage of population receiving social benefits. Daytime LST quartiles were 32.3°C, 33.8°C and 35.0°C.

Figure S3: Difference in mortality increase among those > 65 years of age with a 1°C increase in temperature above the 95th (a) and 99th (b) percentiles lags 0-2 for NDVI quartiles with 95% confidence intervals. GAMs allowing for interaction between temperature and different NDVI classes were adjusted for long-term and seasonal trend (6 df per year), daily averages of O₃ and PM₁₀ (lag 0-1), percent of parish population >65 years of age, building density, proportion of college graduates and percentage of population receiving social benefits.

Figure S4: Difference in mortality increase among those > 65 years of age with a 1°C increase in temperature above the 95th (a) and 99th (b) percentile for lags 0-2 for areas with a distance of > 4 km and ≤ 4 km to the Atlantic Ocean and Tagus Estuary with 95% confidence intervals. GAMs allowing for interaction between temperature and distance classes were adjusted for long-term and seasonal trend (6 df per year), daily averages of O3 and PM10, percent of parish population >65 years of age, building density, proportion of college graduates and percentage of population receiving social benefits.

Figure S5: Difference in mortality increase among those > 65 years of age with a 1°C increase in UTCI above the 95th (a) and 99th (b) percentiles lags 0-2 for nightime LST quartiles with 95% confidence intervals. GAMs allowing for interaction between UTCI and different nighttime LST classes were adjusted for long-term and seasonal trend (6 df per year), daily averages of O₃ and PM₁₀ (lag 0-1), percent of parish population >65 years of age, building density, proportion of college graduates and percentage of population receiving social benefits. Nighttime LST quartiles were 16.6°C, 17.7°C and 18.1°C.

Figure S6: Difference in mortality increase among those > 65 years of age with a 1°C increase in UTCI above the 95th (a) and 99th (b) percentiles lags 0-2 for daytime LST quartiles with 95% confidence intervals. GAMs allowing for interaction between UTCI and different daytime LST classes were adjusted for long-term and seasonal trend (6 df per year), daily averages of O₃ and PM₁₀ (lag 0-1), percent of parish population >65 years of age, building density, proportion of college graduates and percentage of population receiving social benefits. Daytime LST quartiles were 32.3°C, 33.8°C and 35.0°C.